

Important Information About CRESTOR:


Prescribing Information 7 Important Safety Information | Approved Uses |

CRESTOR on the Move Home

Understanding High Cholesterol

Starting with CRESTOR

Save on CRESTOR Stay on Track



Important Safety Information about CRESTOR® (rosuvastatin calcium) Tablets

- CRESTOR is not right for everyone. Do not take CRESTOR if you are nursing, pregnant or may become pregnant; have liver problems; or have had an allergic reaction to CRESTOR
- Your doctor should do blood tests to check your liver before starting treatment with CRESTOR and if you have symptoms of liver problems while taking CRESTOR
- Call your doctor right away if you:

dolor sit amet.

- Have unexplained muscle pain or weakness, especially with fever
- Have muscle problems that do not go away even after your doctor told you to stop taking CRESTOR
- Feel unusually tired
- Have loss of appetite, upper belly pain, dark urine, or yellowing of skin or eyes

These could be signs of rare but serious side effects

- Elevated blood sugar levels have been reported with statins, including CRESTOR
- Side effects: The most common side effects may include headache, muscle aches, abdominal pain, weakness, and nausea. Memory loss and confusion have also been reported with statins, including CRESTOR
- Tell your doctor and pharmacist about other medicines you are taking

Talk to your doctor about prescription CRESTOR.

Approved Uses for CRESTOR

When diet and exercise alone aren't enough to lower cholesterol, adding CRESTOR can help.

In adults, CRESTOR is prescribed along with diet to lower high cholesterol and to slow the buildup of plaque in arteries as part of a treatment plan to lower cholesterol to goal.

Prescribing Information with Patient Information (PDF - 152k)

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.FDA.gov/medwatch or call 1-800-FDA-1088.



This site is intended for US consumers only.

The information on this Web site should not take the place of talking with your doctor or health care professional. If you have any questions about your condition, or if you would like more information about CRESTOR, talk to your doctor or pharmacist. Only you and your health care professional can decide if CRESTOR is right for you.

Legal Statement Contact Us US Corporate Site Privacy Notice Site Map

CRESTOR is licensed from SHIONOGI & CO, LTD, Osaka, Japan.

CRESTOR is a registered trademark and AZ&Me is a trademark of the AstraZeneca group of companies.

LIPITOR is a registered trademark of Pfizer Inc.

©2014 AstraZeneca. All rights reserved. 1733504-2984702 Last Updated 10/14







Important Information About CRESTOR:

Prescribing Information 7 Important Safety Information | Approved Uses |

Home

CRESTOR on the Move


Understanding High Cholesterol

Starting with CRESTOR

Save on CRESTOR

odio convallis.

Stay on Track



Important Safety Information about CRESTOR® (rosuvastatin calcium) Tablets

- CRESTOR is not right for everyone. Do not take CRESTOR if you are nursing, pregnant or may become pregnant; have liver problems; or have had an allergic reaction to CRESTOR
- Your doctor should do blood tests to check your liver before starting treatment with CRESTOR and if you have symptoms of liver problems while taking CRESTOR

odio convallis.

Call your doctor right away if you:

dolor sit amet.

- Have unexplained muscle pain or weakness, especially with fever
- Have muscle problems that do not go away even after your doctor told you to stop taking CRESTOR
- Feel unusually tired
- Have loss of appetite, upper belly pain, dark urine, or yellowing of skin or eyes

These could be signs of rare but serious side effects

- Elevated blood sugar levels have been reported with statins, including CRESTOR
- Side effects: The most common side effects may include headache, muscle aches, abdominal pain, weakness, and nausea. Memory loss and confusion have also been reported with statins, including CRESTOR
- Tell your doctor and pharmacist about other medicines you are taking

Talk to your doctor about prescription CRESTOR.

Approved Uses for CRESTOR

When diet and exercise alone aren't enough to lower cholesterol, adding CRESTOR can help.

In adults, CRESTOR is prescribed along with diet to lower high cholesterol and to slow the buildup of plaque in arteries as part of a treatment plan to lower cholesterol to goal.

Prescribing Information with Patient Information (PDF - 152k)

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.FDA.gov/medwatch or call 1-800-FDA-1088.



This site is intended for US consumers only.

The information on this Web site should not take the place of talking with your doctor or health care professional. If you have any questions about your condition, or if you would like more information about CRESTOR, talk to your doctor or pharmacist. Only you and your health care professional can decide if CRESTOR is right for you.

Legal Statement Contact Us US Corporate Site Privacy Notice Site Map

CRESTOR is licensed from SHIONOGI & CO, LTD, Osaka, Japan.

CRESTOR is a registered trademark and AZ&Me is a trademark of the AstraZeneca group of companies.

LIPITOR is a registered trademark of Pfizer Inc.

©2014 AstraZeneca. All rights reserved. 1733504-2984702 Last Updated 10/14

Important Information About CRESTOR:

Important Safety Information | Approved Uses |

Prescribing Information 7

Home

CRESTOR on the Move

Understanding High Cholesterol

Starting with CRESTOR

Save on CRESTOR

Stay on Track

Events Headline Will Appear Here



Event Title

Location:

Date:

Time:



Event Title

Location:

Date:

Time:



Event Title

Location:

Date:

Booth Details:



Event Title

Location:

Date:

Booth Details:



Photo Headline Will Appear Here















December 2, 2014

Important Safety Information about CRESTOR® (rosuvastatin calcium) Tablets

- · CRESTOR is not right for everyone. Do not take CRESTOR if you are nursing, pregnant or may become pregnant; have liver problems; or have had an allergic reaction to CRESTOR
- Your doctor should do blood tests to check your liver before starting treatment with CRESTOR and if you have symptoms of liver problems while taking CRESTOR
- Call your doctor right away if you:
 - Have unexplained muscle pain or weakness, especially with fever
 - Have muscle problems that do not go away even after your doctor told you to stop taking CRESTOR
 - Feel unusually tired
 - Have loss of appetite, upper belly pain, dark urine, or yellowing of skin or eyes

These could be signs of rare but serious side effects

- Elevated blood sugar levels have been reported with statins, including CRESTOR
- Side effects: The most common side effects may include headache, muscle aches, abdominal pain, weakness, and nausea. Memory loss and confusion have also been reported with statins, including CRESTOR
- Tell your doctor and pharmacist about other medicines you are taking

Talk to your doctor about prescription CRESTOR.

Approved Uses for CRESTOR

When diet and exercise alone aren't enough to lower cholesterol, adding CRESTOR can help.

In adults, CRESTOR is prescribed along with diet to lower high cholesterol and to slow the buildup of plaque in arteries as part of a treatment plan to lower cholesterol to goal.

Prescribing Information with Patient Information (PDF - 152k)

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.FDA.gov/medwatch or call 1-800-FDA-1088.



This site is intended for US consumers only.

The information on this Web site should not take the place of talking with your doctor or health care professional. If you have any questions about your condition, or if you would like more information about CRESTOR, talk to your doctor or pharmacist. Only you and your health care professional can decide if CRESTOR is right for you.

US Corporate Site Privacy Notice Legal Statement Contact Us Site Map

CRESTOR is licensed from SHIONOGI & CO, LTD, Osaka, Japan.

CRESTOR is a registered trademark and AZ&Me is a trademark of the AstraZeneca group of companies.

LIPITOR is a registered trademark of Pfizer Inc.

©2014 AstraZeneca. All rights reserved. 1733504-2984702 Last Updated 10/14